04-17-09




Figure 1. Estimated yearly natural production and in-river escapement of all races of adult Chinook Salmon in the Central Valley rivers and streams. 1952 - 1966 and 1992 - 2008 numbers are calculated in CHINOOKPROD using CDFG Grand Tab in-river escapement data (February 18, 2009). Baseline numbers (1967 - 1991) are from Mills and Fisher (CDFG, 1994).

Source: US Fish and Wildlife Service, accessed from http://www.fws.gov/stockton/afrp/index.asp.




Figure 2. Estimated yearly natural production and in-river escapement of adult fall-run Chinook salmon in the Central Valley rivers and streams. 1952 - 1966 and 1992 - 2008 numbers are from CDFG Grand Tab (February 18, 2009). Baseline numbers (1967 - 1991) are from Mills and Fisher (CDFG, 1994).




Figure 3. Estimated yearly adult natural production, and in river adult escapements of winter-run Chinook salmon in the Central Valley rivers and streams. 1992 - 2008 numbers are from CDFG Grand Tab (February 18, 2009). Baseline numbers (1967-1991) are from Mills and Fisher (CDFG, 1994).




Figure 4. Estimated yearly adult natural production, and in-river adult escapements of late-fall-run Chinook salmon in the Central Valley rivers and streams. 1992 – 2008 numbers are from CDFG Grand Tab (February 18, 2009). Baseline numbers (1967 - 1991) are from Mills and Fisher (CDFG, 1994).




Figure 5. Estimated yearly adult natural production, and in-river adult escapements of spring-run Chinook salmon in the Central Valley rivers and streams. 1960 - 1966 and 1992 - 2008 numbers are from CDFG Grand Tab (February 18, 2009). Baseline numbers (1967 - 1991) are from Mills and Fisher (CDFG, 1994).